

Superintegrable non-autonomous Hamiltonian systems

G. SARDANASHVILY

Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

The Mishchenko–Fomenko theorem on action-angle coordinates for superintegrable autonomous Hamiltonian systems is extended to the non-autonomous ones.

I. INTRODUCTION

The Liouville – Arnold theorem for completely integrable systems^{1–3} and the Mishchenko – Fomenko theorem for the superintegrable ones^{4–6} state the existence of action-angle coordinates around a compact invariant submanifold. These theorems were generalized to the case of non-compact invariant submanifolds.^{7–10} This generalization enable us to analyze completely integrable non-autonomous Hamiltonian systems whose invariant submanifolds are necessarily non-compact.^{11,12} Here we aim to extend this analysis to superintegrable non-autonomous Hamiltonian systems.

We consider a non-autonomous mechanical system whose configuration space is a fibre bundle $Q \rightarrow \mathbb{R}$ over the time axis $\mathbb{R}$ endowed with the Cartesian coordinate t possessing transition functions $t' = t + \text{const}$. Its phase space is the vertical cotangent bundle $V^*Q \rightarrow Q$ of $Q \rightarrow \mathbb{R}$ endowed with the Poisson structure $\{, \}_V$ (13).^{13,14} A Hamiltonian of a non-autonomous mechanical system is a section H (8) of the one-dimensional fibre bundle

$$\zeta : T^*Q \rightarrow V^*Q, \quad (1)$$

where T^*Q is the cotangent bundle of Q endowed with the canonical symplectic form (7).

Definition 1: A non-autonomous Hamiltonian system of $m = \dim Q - 1$ degrees of freedom is called superintegrable if admits $m \leq n < 2m$ integrals of motion $\Phi_1, \dots, \Phi_n$ obeying the following conditions.

(i) All the functions Φ_α are independent, i.e. the n -form $d\Phi_1 \wedge \dots \wedge d\Phi_n$ nowhere vanishes on V^*Q . It follows that the map

$$\Phi : V^*Q \rightarrow N = (\Phi_1(V^*Q), \dots, \Phi_n(V^*Q)) \subset \mathbb{R}^n \quad (2)$$

is a fibred manifold over a connected open subset $N \subset \mathbb{R}^n$.

(ii) There exist smooth real functions s_{ij} on N such that

$$\{\Phi_\alpha, \Phi_\beta\}_V = s_{\alpha\beta} \circ \Phi, \quad \alpha, \beta = 1, \dots, n. \quad (3)$$

(iii) The matrix function with the entries $s_{\alpha\beta}$ (3) is of constant corank $k = 2m - n$ at all points of N .

To describe this non-autonomous Hamiltonian system, we use the fact that there exists an equivalent autonomous Hamiltonian system on the cotangent bundle T^*Q (Theorem 3) which is superintegrable (Theorem 7). Our goal is the following.

Theorem 2: Let Hamiltonian vector fields of the functions Φ_α be complete, and let fibres of the fibred manifold Φ (2) be connected and mutually diffeomorphic. Then there exists an open neighborhood U_M of a fibre M of Φ (2) which is a trivial principal bundle with the structure group

$$\mathbb{R}^{1+k-r} \times T^r \quad (4)$$

whose bundle coordinates are the generalized action-angle coordinates

$$(p_A, q^A, I_\lambda, t, y^\lambda), \quad A = 1, \dots, n - m, \quad \lambda = 1, \dots, k, \quad (5)$$

such that:

- (i) (t, y^λ) are coordinates on the toroidal cylinder (4),
- (ii) the Poisson bracket $\{, \}_V$ on U_M reads

$$\{f, g\}_V = \partial^A f \partial_A g - \partial^A g \partial_A f + \partial^\lambda f \partial_\lambda g - \partial^\lambda g \partial_\lambda f,$$

- (iii) the Hamiltonian H depends only on the action coordinates I_λ ,
- (iv) the integrals of motion $\Phi_1, \dots, \Phi_n$ are independent of coordinates (t, y^λ) .

If $n = m$, we are in the case of a completely integrable non-autonomous Hamiltonian system (Theorem 6).

II. NON-AUTONOMOUS HAMILTONIAN MECHANICS

The configuration space $Q \rightarrow \mathbb{R}$ of a non-autonomous Hamiltonian system is equipped with bundle coordinates (t, q^i) , $i = 1, \dots, m$.^{13,14} Its phase space V^*Q is provided with holonomic coordinates $(t, q^i, p_i = \dot{q}_i)$ with respect to fibre bases $\{\bar{d}q^i\}$ for V^*Q . The cotangent bundle T^*Q of Q plays a role of the homogeneous phase space endowed with holonomic coordinates (t, q^i, p_0, p_i) possessing transition functions

$$p'_i = \frac{\partial q^j}{\partial q'^i} p_j, \quad p'_0 = p_0 + \frac{\partial q^j}{\partial t'} p_j. \quad (6)$$

The cotangent bundle T^*Q admits the canonical symplectic form

$$\Omega = dp_0 \wedge dt + dp_i \wedge dq^i, \quad (7)$$

and the corresponding Poisson bracket

$$\{f, g\} = \partial^0 f \partial_t g - \partial^0 g \partial_t f + \partial^i f \partial_i g - \partial^i g \partial_i f, \quad f, g \in C^\infty(T^*Q).$$

A glance at the transformation law (6) shows that the one-dimensional fibre bundle ζ (1) is a trivial affine bundle. Given its global section

$$H : V^*Q \rightarrow T^*Q, \quad p_0 \circ H = -\mathcal{H}(t, q^j, p_j), \quad (8)$$

the cotangent bundle T^*Q is equipped with the fibre coordinate

$$I_0 = p_0 + \mathcal{H}, \quad I_0 \circ H = 0, \quad (9)$$

possessing the identity transition functions. With respect to the coordinates

$$(t, q^i, I_0, p_i), \quad i = 1, \dots, m, \quad (10)$$

the fibration (1) reads

$$\zeta : \mathbb{R} \times V^*Q \ni (t, q^i, I_0, p_i) \rightarrow (t, q^i, p_i) \in V^*Q. \quad (11)$$

The fibre bundle (1) provides the vertical cotangent bundle V^*Q with the canonical Poisson structure $\{, \}_V$ such that

$$\zeta^* \{f, g\}_V = \{\zeta^* f, \zeta^* g\}, \quad (12)$$

$$\{f, g\}_V = \partial^i f \partial_i g - \partial^i g \partial_i f, \quad f, g \in C^\infty(V^*Q). \quad (13)$$

The Hamiltonian vector fields of functions on V^*Q with respect to the Poisson bracket (13) are vertical vector fields

$$\vartheta_f = \partial^i f \partial_i - \partial_i f \partial^i, \quad f \in C^\infty(V^*Q), \quad (14)$$

$$[\vartheta_f, \vartheta_{f'}] = \vartheta_{\{f, f'\}_V}, \quad (15)$$

on $V^*Q \rightarrow \mathbb{R}$. Accordingly, the corresponding symplectic foliation on the phase space V^*Q coincides with the fibration $V^*Q \rightarrow \mathbb{R}$.

A Hamiltonian of non-autonomous mechanics on the phase space V^*Q is defined as a global section (8) of the affine bundle ζ (1). Then there exists a unique vector field γ_H on V^*Q such that

$$\begin{aligned} \gamma_H \rfloor dt &= 1, & \gamma_H \rfloor H^* \Omega &= 0, \\ \gamma_H &= \partial_t + \partial^i \mathcal{H} \partial_i - \partial_i \mathcal{H} \partial^i. \end{aligned} \quad (16)$$

This vector field, called the Hamilton vector field, defines the first order Hamilton equations

$$q_t^i = \partial^i \mathcal{H}, \quad p_{ti} = -\partial_i \mathcal{H} \quad (17)$$

on a fibre bundle $V^*Q \rightarrow \mathbb{R}$ with respect to the adapted coordinates $(t, q^i, p_i, q_t^i, p_{ti})$ on the first order jet manifold $J^1 V^*Q$ of $V^*Q \rightarrow \mathbb{R}$. Due to the canonical imbedding $J^1 V^*Q \rightarrow TV^*Q$, the Hamilton equations (17) are equivalent to the first order differential equations

$$\dot{t} = 1, \quad \dot{q}^i = \partial^i \mathcal{H}, \quad \dot{p}_i = -\partial_i \mathcal{H} \quad (18)$$

on a manifold V^*Q .

In order to describe evolution of a mechanical system, the Hamilton vector field γ_H (16) is assumed to be complete. It defines a trivialization

$$V^*Q \cong \mathbb{R} \times P$$

which is a canonical automorphism of the Poisson manifold V^*Q such that the corresponding coordinates $(t, \bar{q}^i, \bar{p}_i)$ are the initial date coordinates.¹⁴ With respect to these coordinates, the Hamiltonian (8) reads $\mathcal{H} = 0$, and the Hamilton equations (17) take the form

$$\bar{q}_t^i = 0, \quad \bar{p}_{ti} = 0.$$

We agree to call (V^*Q, H) the non-autonomous Hamiltonian system of m degrees of freedom.

Theorem 3: A non-autonomous Hamiltonian system (V^*Q, H) is equivalent to an autonomous Hamiltonian system $(T^*Q, \mathcal{H}^*)$ of $m + 1$ degrees of freedom on a symplectic manifold (T^*Q, Ω) whose Hamiltonian is the function^{15,16}

$$\mathcal{H}^* = I_0 = p_0 + \mathcal{H}. \quad (19)$$

The Hamiltonian vector field $u_{\mathcal{H}^*}$ of $\mathcal{H}^*$ (19) on T^*Q is

$$u_{\mathcal{H}^*} = \partial_t - \partial_t \mathcal{H} \partial^0 + \partial^i \mathcal{H} \partial_i - \partial_i \mathcal{H} \partial^i. \quad (20)$$

Written relative to the coordinates (10), this vector field reads

$$u_{\mathcal{H}^*} = \partial_t + \partial^i \mathcal{H} \partial_i - \partial_i \mathcal{H} \partial^i. \quad (21)$$

It is projected onto the Hamilton vector field γ_H (16) on V^*Q such that

$$\zeta^*(\mathbf{L}_{\gamma_H} f) = \{\mathcal{H}^*, \zeta^* f\}, \quad f \in C^\infty(V^*Q). \quad (22)$$

The corresponding autonomous Hamilton equations on T^*Q take the form

$$\dot{t} = 1, \quad \dot{p}_0 = -\partial_t \mathcal{H}, \quad \dot{q}^i = \partial^i \mathcal{H}, \quad \dot{p}_i = -\partial_i \mathcal{H}. \quad (23)$$

They are equivalent to the Hamilton equations (18).

Obviously, the vector field $u_{\mathcal{H}^*}$ (21) is complete if the Hamilton vector field γ_H (16) is complete.

III. COMPLETELY INTEGRABLE NON-AUTONOMOUS HAMILTONIAN SYSTEMS

An integral of motion of a non-autonomous Hamiltonian system (V^*Q, H) is defined as a smooth real function F on V^*Q whose Lie derivative

$$\mathbf{L}_{\gamma_H} F = \gamma_H \lrcorner dF = \partial_t F + \{\mathcal{H}, F\}_V$$

along the Hamilton vector field γ_H (16) vanishes. Given the Hamiltonian vector field ϑ_F (14) of F with respect to the Poisson bracket (13), it is easily justified that

$$[\gamma_H, \vartheta_F] = \vartheta_{\mathbf{L}_{\gamma_H} F}. \quad (24)$$

Definition 4: A non-autonomous Hamiltonian system (V^*Q, H) of m degrees of freedom is said to be completely integrable if it admits m integrals of motion $F_1, \dots, F_m$ which are in involution with respect to the Poisson bracket $\{\cdot, \cdot\}_V$ (13) and whose differentials dF_α are linearly independent, i.e.

$$dF_1 \wedge \dots \wedge dF_m \neq 0.$$

By virtue of the relations (15) and (24), the vector fields

$$(\gamma_H, \vartheta_{F_1}, \dots, \vartheta_{F_m}), \quad (25)$$

$$\vartheta_{F_\alpha} = \partial^i F_\alpha \partial_i - \partial_i F_\alpha \partial^i.$$

mutually commute and, therefore, they span an $(m+1)$ -dimensional involutive distribution $\mathcal{V}$ on V^*Q . Let G be the group of local diffeomorphisms of V^*Q generated by the flows of vector fields (25). Maximal integral manifolds of $\mathcal{V}$ are the orbits of G and invariant submanifolds of vector fields (25).^{12,17} They yield a foliation $\mathcal{F}$ of V^*Q .

Let (V^*Q, H) be a non-autonomous Hamiltonian system and $(T^*Q, \mathcal{H}^*)$ an equivalent autonomous Hamiltonian system on T^*Q . An immediate consequence of the relations (12) and (22) is the following.⁹

Theorem 5: Given a completely integrable non-autonomous Hamiltonian system

$$(\gamma_H, F_1, \dots, F_m) \quad (26)$$

of m degrees of freedom on V^*Q , the autonomous Hamiltonian system

$$(\mathcal{H}^*, \zeta^* F_1, \dots, \zeta^* F_m) \quad (27)$$

of $m + 1$ degrees of freedom on T^*Q is completely integrable.

The Hamiltonian vector fields

$$\begin{aligned} & (u_{\mathcal{H}^*}, u_{\zeta^* F_1}, \dots, u_{\zeta^* F_m}), \\ & u_{\zeta^* F_\alpha} = \partial^i F_\alpha \partial_i - \partial_i F_\alpha \partial^i, \end{aligned} \quad (28)$$

of the autonomous integrals of motion (27) span an $(m + 1)$ -dimensional involutive distribution $\mathcal{V}_T$ on T^*Q such that

$$T\zeta(\mathcal{V}_T) = \mathcal{V}, \quad TH(\mathcal{V}) = \mathcal{V}_T|_{H(V^*Q)=I_0=0}, \quad (29)$$

where

$$TH : TV^*Q \ni (t, q^i, p_i, \dot{t}, \dot{q}^i, \dot{p}_i) \rightarrow (t, q^i, p_i, I_0 = 0, \dot{t}, \dot{q}^i, \dot{p}_i, \dot{I}_0 = 0) \in TT^*Q.$$

It follows that, if M is an invariant submanifold of the completely integrable non-autonomous Hamiltonian system (26), then $H(M)$ is an invariant submanifold of the completely integrable autonomous Hamiltonian system (27).

In order to introduce generalized action-angle coordinates around an invariant submanifold M of the completely integrable non-autonomous Hamiltonian system (26), let us suppose that the vector fields (25) on M are complete. It follows that M is a locally affine manifold diffeomorphic to a toroidal cylinder

$$(\mathbb{R}^{1+m-r} \times T^r). \quad (30)$$

Moreover, let us assume that there exists an open neighbourhood U of M such that the foliation $\mathcal{F}$ of U is a fibred manifold $\phi : U \rightarrow N$ over a domain $N \subset \mathbb{R}^m$ whose fibres are mutually diffeomorphic.¹¹

Because the morphism TH (29) is a bundle isomorphism, the Hamiltonian vector fields (28) on the invariant submanifold $H(M)$ of the completely integrable autonomous Hamiltonian system are complete. Since the affine bundle ζ (11) is trivial, the open neighbourhood $\zeta^{-1}(U)$ of the invariant submanifold $H(M)$ is a fibred manifold

$$\tilde{\phi} : \zeta^{-1}(U) = \mathbb{R} \times U \xrightarrow{(\text{Id}, \phi)} \mathbb{R} \times N = N'$$

over a domain $N' \subset \mathbb{R}^{m+1}$ whose fibres are diffeomorphic to the toroidal cylinder (30). In accordance with the Liouville – Arnold theorem extended to the case of non-compact invariant submanifolds,⁷ the open neighbourhood $\zeta^{-1}(U)$ of $H(M)$ is a trivial principal bundle

$$\zeta^{-1}(U) = N' \times (\mathbb{R}^{1+m-r} \times T^r) \rightarrow N' \quad (31)$$

with the structure group (30) whose bundle coordinates are the generalized action-angle coordinates

$$(I_0, I_1, \dots, I_m, t, z^1, \dots, z^m) \quad (32)$$

such that:

- (i) (t, z^a) are coordinates on the toroidal cylinder (30),
- (ii) the symplectic form (7) on $\zeta^{-1}(U)$ reads

$$\Omega = dI_0 \wedge dt + dI_a \wedge dz^a,$$

- (iii) $\mathcal{H}^* = I_0$,

- (iv) the integrals of motion $\zeta^* F_1, \dots, \zeta^* F_m$ depend only on the action coordinates $I_1, \dots, I_m$.

Provided with the coordinates (32), $\zeta^{-1}(U) = U \times \mathbb{R}$ is a trivial bundle possessing the fibre coordinate I_0 (9). Consequently, the open neighbourhood U of an invariant submanifold M of the completely integrable non-autonomous Hamiltonian system (25) is diffeomorphic to the Poisson annulus

$$U = N \times (\mathbb{R}^{1+m-r} \times T^r) \quad (33)$$

endowed with the generalized action-angle coordinates

$$(I_1, \dots, I_m, t, z^1, \dots, z^m) \quad (34)$$

such that:

- (i) the Poisson structure (13) on U takes the form

$$\{f, g\}_V = \partial^a f \partial_a g - \partial_a g \partial_a f,$$

- (ii) the Hamiltonian (8) reads $\mathcal{H} = 0$,

- (iii) the integrals of motion $F_1, \dots, F_m$ depend only on the action coordinates $I_1, \dots, I_m$.

The Hamilton equations (17) relative to the generalized action-angle coordinates (34) take the form

$$z_t^a = 0, \quad I_{ta} = 0.$$

It follows that the generalized action-angle coordinates (34) are the initial date coordinates.

Note that the generalized action-angle coordinates (34) by no means are unique. Given a smooth function $\mathcal{H}'$ on $\mathbb{R}^m$, one can provide $\zeta^{-1}(U)$ with the generalized action-angle coordinates

$$t, \quad z'^a = z^a - t\partial^a \mathcal{H}', \quad I'_0 = I_0 + \mathcal{H}'(I_b), \quad I'_a = I_a.$$

With respect to these coordinates, a Hamiltonian of the autonomous Hamiltonian system on $\zeta^{-1}(U)$ reads $\mathcal{H}'^* = I'_0 - \mathcal{H}'$. A Hamiltonian of the non-autonomous Hamiltonian system on U endowed with the generalized action-angle coordinates (t, z'^a, I_a) is $\mathcal{H}'$.

Thus, the following has been proved.

Theorem 6: Let $(\gamma_H, F_1, \dots, F_m)$ be a completely integrable non-autonomous Hamiltonian system. Let M be its invariant submanifold such that the vector fields (25) on M are complete and there exists an open neighbourhood U of M which is a fibred manifold in mutually diffeomorphic invariant submanifolds. Then U is diffeomorphic to the Poisson annulus (33), and it can be provided with the generalized action-angle coordinates (34) such that the integrals of motion $(F_1, \dots, F_m)$ and the Hamiltonian H depend only on the action coordinates $I_1, \dots, I_m$.

IV. SUPERINTEGRABLE NON-AUTONOMOUS HAMILTONIAN SYSTEMS

Let $(\gamma_H, \Phi_1, \dots, \Phi_n)$ be a superintegrable non-autonomous Hamiltonian system in accordance with Definition 1. The associated autonomous Hamiltonian system on T^*Q possesses $n + 1$ integrals of motion

$$(\mathcal{H}^*, \zeta^* \Phi_1, \dots, \zeta^* \Phi_n) \tag{35}$$

with the following properties.

(i) The functions (35) are mutually independent, and the map

$$\begin{aligned} \tilde{\Phi} : T^*Q &\rightarrow (\mathcal{H}^*(T^*Q), \zeta^* \Phi_1(T^*Q), \dots, \zeta^* \Phi_n(T^*Q)) = \\ &(I_0, \Phi_1(V^*Q), \dots, \Phi_n(V^*Q)) = \mathbb{R} \times N = N' \end{aligned} \tag{36}$$

is a fibred manifold.

(ii) The functions (35) obey the relations

$$\{\zeta^* \Phi_\alpha, \zeta^* \Phi_\beta\} = s_{\alpha\beta} \circ \zeta^* \Phi, \quad \{\mathcal{H}^*, \zeta^* \Phi_\alpha\} = s_{0\alpha} = 0$$

so that the matrix function with the entries $(s_{0\alpha}, s_{\alpha\beta})$ on N' is of constant corank $2m + 1 - n$.

Refereing to the definition of an autonomous superintegrable system,⁴⁻⁶ we come to the following.

Theorem 7: Given a superintegrable non-autonomous Hamiltonian system (γ_H, Φ_α) on V^*Q , the associated autonomous Hamiltonian system (35) on T^*Q is superintegrable.

There is the commutative diagram

$$\begin{array}{ccc} T^*Q & \xrightarrow{\zeta} & V^*Q \\ \tilde{\Phi} \downarrow & & \downarrow \Phi \\ N' & \xrightarrow{\xi} & N \end{array}$$

where ζ (11) and

$$\xi : N' = \mathbb{R} \times N \rightarrow N$$

are trivial bundles. It follows that the fibred manifold (36) is the pull-back $\tilde{\Phi} = \xi^*\Phi$ of the fibred manifold Φ (2) onto N' .

Let the conditions of Theorem 2 hold. If the Hamiltonian vector fields

$$(\gamma_H, \vartheta_{\Phi_1}, \dots, \vartheta_{\Phi_n}), \quad \vartheta_{\Phi_\alpha} = \partial^i \Phi_\alpha \partial_i - \partial_i \Phi_\alpha \partial^i,$$

of integrals of motion Φ_α on V^*Q are complete, the Hamiltonian vector fields

$$(u_{\mathcal{H}^*}, u_{\zeta^*\Phi_1}, \dots, u_{\zeta^*\Phi_n}), \quad u_{\zeta^*\Phi_\alpha} = \partial^i \Phi_\alpha \partial_i - \partial_i \Phi_\alpha \partial^i,$$

on T^*Q are complete. If fibres of the fibred manifold Φ (2) are connected and mutually diffeomorphic, the fibres of the fibred manifold $\tilde{\Phi}$ (36) also are well.

Let M be a fibre of Φ (2) and $H(M)$ the corresponding fibre of $\tilde{\Phi}$ (36). In accordance with the Mishchenko – Fomenko theorem extended to the case of non-compact invariant submanifolds,^{9,10} there exists an open neighbourhood U' of $H(M)$ which is a trivial principal bundle with the structure group (4) whose bundle coordinates are the generalized action-angle coordinates

$$(p_A, q^A, I_0, I_\lambda, t, y^\lambda), \quad A = 1, \dots, n - m, \quad \lambda = 1, \dots, k, \quad (37)$$

such that:

- (i) (t, y^λ) are coordinates on the toroidal cylinder (4),
- (ii) the symplectic form Ω (7) on U' reads

$$\Omega = dp_A \wedge dq^A + dI_0 \wedge dt + dI_\alpha \wedge dy^\alpha,$$

(iii) the action coordinates (I_0, I_α) are expressed into the values of the Casimir functions $C_0 = I_0$, C_α of the coinduced Poisson structure

$$w = \partial^A \wedge \partial_A$$

on N' ,

- (iv) the Hamiltonian $\mathcal{H}^*$ depends on the action coordinates, namely, $\mathcal{H}^* = I_0$,
- (iv) the integrals of motion $\zeta^*\Phi_1, \dots, \zeta^*\Phi_n$ are independent of the coordinates (t, y^λ) .

Provided with the generalized action-angle coordinates (37), the above mentioned neighbourhood U' is a trivial bundle $U' = \mathbb{R} \times U_M$ where $U_M = \zeta(U')$ is an open neighbourhood of the fibre M of the fibre bundle Φ (2). As a result, we come to Theorem 2.

- ¹ V.Arnold and A.Avez, *Ergodic Problems in Classical Mechanics* (Benjamin, New York, 1968).
- ² V.Arnold (Ed.), *Dynamical Systems III, IV* (Springer-Verlag, Berlin, 1990).
- ³ V.Lazutkin, *KAM Theory and Semiclassical Approximations to Eigenfunctions* (Springer-Verlag, Berlin, 1993).
- ⁴ A.Mishchenko and A.Fomenko, *Funct. Anal. Appl.* **12**, 113 (1978).
- ⁵ A.Bolsinov and B.Jovanović, *Ann. Global Anal. Geom.* **23**, 305 (2003).
- ⁶ F.Fassó, *Acta Appl. Math.* **87**, 93 (2005).
- ⁷ E.Fiorani, G.Giachetta and G.Sardanashvily, *J. Phys. A* **36**, L101 (2003); *E-print arXiv: math.DS/0210346*.
- ⁸ G.Giachetta, L.Mangiarotti and G.Sardanashvily, *J. Math. Phys.* **44**, 1984 (2003); *E-print arXiv math.DS/0211463*.
- ⁹ E.Fiorani and G.Sardanashvily, *J. Phys. A* **39**, 14035 (2006); *E-print arXiv: math.DG/0604104*.
- ¹⁰ E.Fiorani and G.Sardanashvily, *J. Math. Phys.* **48**, 032001 (2007); *E-print arXiv: math.DG/0610790*.
- ¹¹ G.Giachetta, L.Mangiarotti and G.Sardanashvily, *J. Phys. A* **35**, L439 (2002); *E-print arXiv: math.DS/0204151*.
- ¹² G.Giachetta, L.Mangiarotti and G.Sardanashvily, *Geometric and Algebraic Topological Methods in Quantum Mechanics* (World Scientific, Singapore, 2005).

- ¹³ G.Sardanashvily, J. Math. Phys. **39**, 2714 (1998).
- ¹⁴ L.Mangiarotti and G.Sardanashvily, *Gauge Mechanics* (World Scientific, Singapore, 1998).
- ¹⁵ A.Dewismw and S.Bouquet, J. Math. Phys. **34**, 997 (1993).
- ¹⁶ L.Mangiarotti and G.Sardanashvily, J. Math. Phys. **41**, 2858 (2000).
- ¹⁷ H. Sussmannn, Trans. Amer. Math. Soc. **180**, 171 (1973).